

Inelastic Scattering of Tritium-Source Antineutrinos on Electrons of Germanium Atoms

V. I. Kopeikin, L. A. Mikaelyan, and V. V. Sinev

Russian Research Centre Kurchatov Institute, Moscow, Russia

Processes of the inelastic magnetic and weak scattering of tritium-beta-source antineutrinos on the bound electrons of a germanium atom are considered. The results obtained by calculating the spectra and cross sections are presented for the energy-transfer range between 1 eV and 18 keV.

Introduction

This study was performed in connection with the experiment planned by researchers from three institutions (Institute of Theoretical and Experimental Physics, Joint Institute for Nuclear Research, and the All-Russia Research Institute for Experimental Physics) to look for the magnetic moment of the electron neutrino [1]. This experiment is presumed to study the spectra of energy deposited in a detector material upon the inelastic scattering of tritium-source antineutrinos on atomic electrons. The sensitivity to the magnitude of the neutrino magnetic moment is assumed to be $\mu \sim 3 \times 10^{-12} \mu_B$ (μ_B is the Bohr magneton), which is nearly two orders of magnitude better than the upper limit, $\mu \leq 2 \times 10^{-10} \mu_B$, set in experiments performed at the Savannah River nuclear reactors, the Rovensk nuclear power plant, and Krasnoyarsk neutrino laboratory [2].

The discovery of the neutrino magnetic moment would suggest the existence of phenomena beyond the fundamentals of electroweak-interaction theory and would have far-reaching consequences for particles physics and astrophysics. (For a discussion on the issues concerning the motivation of searches for the neutrino magnetic moment in laboratory experiments and the status and prospects of investigations along these lines, see, for example, the original study of Neganov *et al.* [1], the review articles quoted in [3], and references therein.)

The magnetic moment can be sought by analyzing the measured spectrum of events of $\bar{\nu}_e e$ scattering. In the case of scattering on a free electron, the cross section for weak $\bar{\nu}_e e$ scattering tends to a finite limit, while the cross

section for magnetic scattering grows indefinitely. Therefore, the experimental sensitivity to the magnitude of the neutrino magnetic moment increases with decreasing energy of measured events. As the energy of an event decreases, however, the effects of electron binding in target atoms begin to affect relevant cross sections and their energy dependence.

A feature peculiar to the experiment proposed in [1] is that, in the majority of cases, the energy q deposited in the target upon the scattering of tritium neutrinos $\bar{\nu}_e$ is below 1 keV. The experiment will employ germanium and silicon detectors that, owing to the internal-signal-amplification mechanism, will make it possible to perform measurements in the region of energies as low as those indicated immediately above.

In this study, we consider the inelastic scattering of tritium-source antineutrinos on germanium-atom electrons. The ensuing exposition is organized as follows. First, we recall the expression for the cross sections for elastic $\bar{\nu}_e e$ scattering, whereupon we briefly dwell on features of inelastic scattering and on the procedure for calculating relevant cross sections. Further, we display the results obtained by calculating the spectra of inelastic scattering on electrons of germanium-atom subshells and, finally, present the total spectra and integrated cross sections in a form that is suitable for a direct comparison with experimental data.

1 Elastic and inelastic scattering of antineutrinos on an electron

1. The differential cross section for the magnetic scattering of a neutrino on a free electron at rest is proportional to the squared magnetic moment μ^2 [4]; that is,

$$(d\sigma^M/dq)_{free} = \pi r_0^2 (\mu/\mu_B)^2 (1/q - 1/E), \quad (1)$$

where $\pi r_0^2 = 2.495 \times 10^{-25} \text{ cm}^2$; E is the incident-neutrino energy; and q is the energy transfer, which coincides, in this case, with the recoil-electron kinetic energy.

The differential cross section for the electroweak scattering of an antineutrino on a free electron at rest has the form (see, for example, [5])

$$(d\sigma^W/dq)_{free} = G_F^2 (m/2\pi) [4x^4 +$$

$$+(1+2x^2)^2(1-q/E)^2-2x^2(1+2x^2)mq/E^2], \quad (2)$$

where m is the electron mass; $G_F^2(m/2\pi) = 4.31 \times 10^{-48} \text{ cm}^2\text{keV}$; and $x^2 = \sin^2 \theta_W = 0.232$, θ_W being the Weinberg angle.

For a given value of primary neutrino energy E , the energy transfer to the electron involved is restricted by the kinematical condition

$$q \leq q_{max} = 2E^2/(2E + m). \quad (3)$$

In the region $q > q_{max}$, free-scattering cross sections are zero. In the case of tritium-source antineutrinos $\bar{\nu}_e$ whose endpoint energy is 18.6 keV, $q_{jmax} = 1.26 \text{ keV}$.

With decreasing energy, the cross section for free magnetic scattering increases in proportion to $1/q$, while the cross section (2) tends to a finite limit of $1.016 \times 10^{-47} \text{ cm}^2\text{keV}$. At the magnetic-moment value of $\mu = 3 \times 10^{-12}$, the spectra of weak and magnetic antineutrino ($\bar{\nu}_e$) scattering intersect at an energy transfer of $q \approx 0.3 \text{ keV}$.

2. In calculating the differential cross sections $d\sigma_i^M/dq$ and $d\sigma_i^W/dq$ for inelastic scattering on the electrons of the i -th shell, we use the approach developed in [6]. In the initial state, there are a germanium atom ($Z = 32$) and incident antineutrinos of energy E ; in the final state, there are a knock-on electron, an outgoing neutrino, and a vacancy in one of the atomic shells. The wave functions for the target atom and the electron binding energies are calculated in the self-consistent relativistic Hartree-Fock-Dirac model. The wave functions for the knock-on electron can be found by numerically integrating the Dirac equation in the same self-consistent potential. The energy-transfer spectra $S_i^M(q)$ and $S_i^W(q)$ are obtained as convolutions of the differential cross sections $d\sigma_i^M/dq$

Table 1: Calculated electron binding energies ϵ_i (in keV) in germanium atomic shells (n_i , is the number of electrons in a shell)

	K ($1s_{1/2}$)	L_I ($2s_{1/2}$)	L_{II} ($2p_{1/2}$)	L_{III} ($2p_{3/2}$)	M_I ($3s_{1/2}$)	M_{II} ($3p_{1/2}$)	M_{III} ($3s_{3/2}$)	M_{IV} ($3d_{2/2}$)	M_V ($3d_{5/2}$)	N_I ($4s_{1/2}$)	N_{II} ($4p_{1/2}$)
n_i	2	2	2	4	2	2	4	4	6	2	2
ϵ_i	10.9	1.37	1.22	1.19	0.17	0.12	0.115	0.031	0.03	(0.013)	(0.005)

and $d\sigma_i^W/dq$ with the spectrum $\rho(E)$ of the tritium-source antineutrinos (Fig. 1) that was calculated by using the Coulomb function [7]. The resulting spectra $S^{M,W}(q)$ for scattering on a germanium atom can be derived by taking

a sum over the shells of the spectra $S_i^{M,W}$ with allowance for the number n_i , of electrons in each shell:

$$S^{M,W}(q) = \sum (n_i/Z) S_i^{M,W}(q). \quad (4)$$

All the spectra $S_i^{M,W}$ and the resulting spectra $S^{M,W}(q)$ are normalized to one electron and are calculated in units of $10^{-47} \text{ cm}^2/(\text{keV electron})$. In all the calculations of cross sections and magnetic-scattering spectra, the neutrino magnetic moment is taken to be $5 \times 10^{-12} \mu_B$. The energy transfer q to the atom involved in an inelastic-scattering event is the sum of the kinetic energy T of the knock-on electron and the electron binding energy ϵ_i , in the i th atomic shell:

$$q = T + \epsilon_i. \quad (5)$$

The radiation generated upon the filling of the vacancy and the knock-on electron are absorbed in a detector material. As a result, the value determined experimentally for the energy of an event coincides with the energy transfer q in this collision event.

The cross sections for weak and magnetic inelastic scattering on an electron of the i th shell vanish for $q \leq \epsilon_i$. (We disregard transitions of the knock-on electron to discrete excited optical levels of the target atom.) In contrast to what is known for free scattering, these cross sections do not vanish at $q = q_{max}$ [see (3)]; in the "forbidden" region $q > q_{max}$, the inelastic-scattering cross sections decrease and vanish only at the q value equal to the incident-antineutrino energy E .

The calculated electron binding energies in the shells of a free germanium atom (Table 1) agree with experimental data to within 5%. This is not true only for four valence electrons, but their binding energy is so low that, in calculating cross sections and spectra, we use, for them, the formulas for scattering on free electrons.

2 Discussion of the results for spectra and cross sections

The scattering-event spectra $S_i^{M,W}(q)$ depends substantially on the binding energy ϵ_i , in a shell, their shapes being different in the regions $q > q_{max}$ and $q < q_{max}$, where $q_{max} = 1 - 26 \text{ keV}$ is the kinematical limit given by (3).

Table 2: Integrated cross sections $I^M(q)$ and $I^W(q)$ (in units of 10^{-47} cm²/electron) for the inelastic magnetic ($\mu = 5 \times 10^{-12}$) and weak scattering of tritium neutrinos on germanium-atom electrons $\bar{\nu}_e e$ in the energy range 0.01- q keV

q , keV	$I^M(q)$	$I^W(q)$
0.02	0.054	0.0012
0.03	0.0855	0.0025
0.05	0.1372	0.0061
0.1	0.2386	0.0173
0.2	0.4046	0.0455
0.35	0.5733	0.0892
0.5	0.6771	0.1254
0.75	0.7225	0.1671
1.	0.8184	0.1907
1.26	0.842	0.2031
2.03	0.8771	0.2186
3.	0.887	0.2228
5.	0.8903	0.2241
10.94	0.8908	0.2243
18.6	0.8909	0.2243

Figures 2 and 3 display the results obtained by calculating the spectra of inelastic scattering on electrons of the L and M germanium shells in the energy-transfer range $q = 1-4$ keV, whose larger part belongs to the region $q > q_{max}$, which is forbidden for free scattering. With the exception of a small local enhancement near the threshold for the scattering of antineutrinos $n\bar{\nu}_e e$ on electrons of the L shells, we observe a decrease in spectra with increasing energy q . A general trend is that the lower the electron binding energy in a shell, the steeper this decrease.

In the energy region $q < q_{max}$ (Figs. 4, 5), L -shell electrons make virtually no contribution. This region is dominated by the scattering on electrons of M atomic shells and on valence electrons. The spectra of magnetic scattering on M shells in the immediate vicinity of the threshold lie much lower than the spectrum of scattering on a free electron. With increasing energy, the distinction between these spectra decreases fast; in a certain energy interval, they virtually coincide, while, for $q > 0.8$ keV, the inelastic-scattering spectra

lie above the free-scattering spectrum. A similar behavior is also observed for weak scattering, but the spectra of elastic and inelastic scattering approach each other more slowly in this case.

The spectra $S^M(q)$ and $S^W(q)$ of magnetic and weak inelastic scattering that include the contributions from all germanium atomic shells are represented by the solid curves in Figs. 6 and 7. They can be used to perform a comparison with experimental data. The numbers of scattering events occurring within a given energy interval are determined by the integrals $I^M(q)$ and $I^W(q)$ of the spectra $S^M(q)$ and $S^W(q)$. Figure 8 and Table 2 display the integrals from the lower detection threshold of 0.01 keV to a variable upper boundary q .

From the data given in Figs. 7 and 8 and in Table 2, it can be seen that, for the chosen value of the neutrino magnetic moment, the magnetic scattering is more intense than weak scattering over the entire energy region. It should be noted that nearly 80% of all magnetic-scattering events fall within the energy-transfer range from 0.05 to 1.26 keV; the energy region of $q > q_{max}$ involves only about 6% of such scattering events; and their fraction in the interval from 0.01 to 0.05 keV amounts to about 15%.

It is interesting to compare the "exact" spectra $S^{M,W}(q)$ and integrated cross sections $I^{M,W}(q)$ with the values obtained in the step-function approximation. In this approximation, it is assumed that the spectrum of inelastic scattering on an electron of the i th shell is coincident with the spectrum of scattering on a free electron for energies of $q > \epsilon_i$, and that it vanishes for $q < \epsilon_i$. In this approximation, the "exact" expression (4) for the resulting spectra reduces to the form

$$S_{SF}^{M,W}(q) = S_{free}^{M,W}(q) \sum (n_i/Z) \theta(q - \epsilon_i). \quad (6)$$

where summation is performed over all atomic shells and $\theta(q - \epsilon_i)$ is the Heaviside step function equal to unity for $q > \epsilon_i$, and to zero for $q < \epsilon_i$. It was shown in [8] that, in the case where antineutrinos generated by a reactor and by a ^{90}Sr - ^{90}Y source are scattered on electrons of iodine and germanium atoms, the approximate spectra (6) agree with the spectra in (4) to within 2% for the energy transfer q ranging from 1-1.5 to 2-3 keV.

From the data shown in Fig. 6, it can be seen that the approximation specified by Eq. (6) is inadequate for both magnetic and weak scattering in the case of a soft spectrum of tritium antineutrinos. The integrated cross sections found in the step-function approximation exceed the corresponding "exact" values by about 25%(Fig. 8).

ACKNOWLEDGMENTS

We would like to emphasize the leading role of S.A. Fayans, who prematurely passed away in 2001, in developing and applying the procedure used in this study. We are grateful to L.N. Bogdanova and A.S. Starostin for stimulating discussions. This work was supported by the Russian Foundation for Basic Research; by the program Leading Scientific Schools (grant No. 00-15-96708); and, in part, by the Department of Atomic Science and Technology of the Ministry for Atomic Industry of the Russian Federation (contract No. 6.25.19.19.02.969).

REFERENCES

1. B. S. Neganove *et al.*, *Yad. Fiz.* 64, 2033 (2001) [*Phys. At.Nucl.*64, 1948(2001)].
2. F. Reines, H. Gurr, and H. Sobel, *Phys. Rev. Lett.* 37, 315 (1976); G. S. Vidyakin, V. N. Vyrodov, I. I. Gurevich *et al.*, *Pis'ma Zh. Eksp. Teor. Fiz.* 55, 212 (1992) [*JETP Lett.* 55,206 (1992)]; A. V. Derbin *et al.*, *Pis'ma Zh. Eksp. Teor. Fiz.* 57,755 (1993) [*JETP Lett.* 57,768 (1993)].
3. A. V. Derbin, *Yad. Fiz.* 57, 236 (1994) [*Phys. At. Nucl.* 57, 222 (1994)]; L. A. Mikaelyan, *Yad. Fiz.* 65, 1206 (2002) [*Phys. At. Nucl.* 65, 1173 (2002)].
4. H. Bethe, *Proc. Cambridge Philos. Soc.* 35, 108 (1935); G. V. DomogatskiT and D. K. Nadezhin, *Yad. Fiz.* 12, 1233 (1970) [*Sov J. Nucl. Phys.* 12, 678 (1971)].
5. M. B. Voloshin, M. I. VysotskiT, and L. B. Okun', *Zh. Eksp. Teor. Fiz.* 91, 754 (1986) [*Sov. Phys. JETP* 64, 446(1986)].
6. S.A. Fayans, V. Yu. Dobretsov, and A. B. Dobrotsvetov, *Phys. Lett. B* 291, 1 (1992); V. Yu. Dobretsov, A. B. Dobrotsvetov, and S. A. Fayans, *Yad. Fiz.* 55, 2126 (1992) [*SovJ.Nucl.Phys.*55, 118 (1992)].
7. B. S. Dzhelepov *et al.* *Beta Processes* (Nauka, Leningrad, 1972).
8. S.A. Fayans, L. A. Mikaelyan, and V. V. Sinev, *Yad. Fiz.* 64, 1551 (2001) [*Phys. At. Nucl.* 64, 1475 (2001)]; V. I. Kopeikin *et al.*, *Yad. Fiz.* 60, 2032 (1997) [*Phys. At.Nucl.*60, 1859(1997)].




Fig. 1. Spectrum $\rho(E)$ of tritium-source antineutrinos.




Fig. 2. Spectra $S_i^M(q)$ of the inelastic magnetic ($\mu = 5 \times 10^{-12} \mu_B$) scattering of antineutrinos on electrons of the L_{I} , L_{III} , M_{I} , M_{III} , M_{V} germanium-atom shells in the energy-transfer range $q = 1-4$ keV. The dashed curve corresponds to scattering on a free electron.




Fig. 3. As in Fig. 2, but for the inelastic weak scattering of $\bar{\nu}_e$.




Fig. 4. Spectra $S_i^M(q)$ of the inelastic magnetic ($\mu = 5 \times 10^{-12} \mu_B$) scattering of antineutrinos on electrons of the L_I , L_{III} , M_I , M_{III} , M_V germanium-atom shells in the energy-transfer range $q = 0.01-1.5$ keV. The dashed curve corresponds to scattering on a free electron.




Fig. 5. As in Fig. 4, but for the inelastic weak scattering of $\bar{\nu}_e$.




Fig. 6. Spectra of the inelastic magnetic and weak antineutrino scattering [$S^M(q)$ ($\mu = 5 \times 10^{-12} \mu_B$) and $S^W(q)$, respectively] on germanium-atom electrons in the energy-transfer range of $q = 0.01$ – 1.5 keV. The dashed curves correspond to the step-function approximation.




Fig. 7. As in Fig. 6, but in the energy-transfer range $q = 1\text{--}4$ keV.




Fig. 8. Integrated cross sections $I^{M,W}(q)$ for magnetic ($\mu = 5 \times 10^{-12} \mu_B$) and weak scattering of antineutrinos on germanium-atom electrons versus the upper boundary of the energy transfer q . The dashed curves correspond to the step-function approximation.